PerCP/Cyanine5.5 anti-mouse CD366 (Tim-3)

Catalog # / 1198585 / 25 μg

Size: 1198590 / 100 μg

Clone: RMT3-23

Isotype: Rat IgG2a, κ

Reactivity: Mouse

Preparation: The antibody was purified by affinity


chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cyanine5.5 and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.5



Mouse Tim-3 transfected cells were stained with anti-mouse CD366 (Tim-3, clone RMT3-23) PerCP/Cy5.5 (filled histogram) or rat IgG2a, κ PerCP/Cy5.5 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal

performance for each application.

* PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum

emission of 690 nm.

Application Notes:

Additional reported applications (for relevant formats) include: *in vitro*1 and *in vitro*2 blocking of Tim 2, and immunohistochemical staining of frazen

vivo2 blocking of Tim-3, and immunohistochemical staining of frozen

µm filtered) is recommended for functional assays (Cat. No. 119707).

sections2. The LEAF™ purified antibody (Endotoxin EU/microg, Azide-Free, 0.2

Application References:

1. Sabatos CA, et al. 2003. Nat. Immunol. 4:1102.

2. Sanchez-Fueyo A, et al. 2003. Nat. Immunol. 4:1102.

3. Kuchroo VK, et al. 2003. Nat. Rev. Immunol. 3:454.

4. Mooney L, et al. 2002. Nature 415:536.

Description: CD366 (Tim-3) is a transmembrane protein also known as T cell

immunoglobulin and mucin domain containing protein-3. Tim-3 is expressed at high levels on Th1 lymphocytes and CD11b⁺ macrophages. Tim-3 has also been shown to exist as a soluble protein. Cells expressing Tim-3 are present at high levels in the CNS of animals at the onset of experimental autoimmune encephalomyelitis (EAE), a disease mediated by lymphocytes secreting Th1-like cytokines. Tim-3 has been proposed to inhibit Th1-mediated immune

responses and promote immunological tolerance.

Antigen References:

- 1. Sabatos CA, et al. 2003. Nat. Immunol. 4:1102.
- 2. Sanchez-Fueyo A, et al. 2003. Nat. Immunol. 4:1102.
- 3. Kuchroo VK, et al. 2003. Nat. Rev. Immunol. 3:454.
- 4. Mooney L, et al. 2002. Nature 415:536.