

PE anti-mouse MAdCAM-1

Catalog # / Size: 1203545 / 25 µg
1203550 / 100 µg

Clone: MECA-367

Isotype: Rat IgG2a, κ


Immunogen: Endothelial cells

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: Lot-specific



TNFα-stimulated bEND.3 cells were stained with anti-mouse MAdCAM-1 (clone MECA-367) PE (filled histogram) or mouse IgG2a, κ isotype control PE (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: *in vitro* and *in vivo* blocking of lymphocyte adhesion and *in vivo* blocking of lymphocyte homing^{1-4,7}, immunohistochemical staining^{1,5,6} of acetone-fixed frozen sections, immunoprecipitation, and Western blotting¹.

- Application References:**
1. Streeter PR, *et al.* 1988. *Nature* 331:41.
 2. Briskin MJ, *et al.* 1993. *Nature* 363:461.
 3. Berlin C, *et al.* 1993. *Cell* 74:185.
 4. Bargatze RF, *et al.* 1995. *Immunity* 3:99.
 5. Tanneau GM, *et al.* 1999. *J. Histochem. Cytochem.* 47:1581.
 6. Savinov AY, *et al.* 2003. *J. Exp. Med.* 197:643.
 7. Rivera-Nieves J, *et al.* 2005. *J. Immunol.* 174:2343.
 8. Hindley JP, *et al.* 2012. *Cancer Res.* 72:5473. [PubMed](#).

Description: MAdCAM-1 is a 58-66kD type I glycoprotein, also known as Mucosal addressin cell adhesion molecule-1. This mucosal vascular addressin is a member of the Ig superfamily found on fetus and neonatal endothelial cells. In adults, MAdCAM-1 is

predominately expressed on high endothelial venule (HEV) of Peyer's patches, mesenteric lymph nodes and gut lamina propria. It is also expressed on vascular endothelium in mammary glands and pancreas. MAdCAM-1, through its interaction with integrin $\alpha 4\beta 7$ or CD62L, is involved in lymphocyte tethering, rolling, and homing. It has been reported that immobilized MAdCAM-1 is able to co-stimulate T cell proliferation. The MECA-367 antibody blocks the interaction of MAdCAM-1 with its counter-receptor both *in vitro* and *in vivo*. *In vivo* administration of the mAb is able to reduce T-cell mediated inflammation in some gastrointestinal diseases.

Antigen
References:

1. Streeter PR, *et al.* 1988. *Nature* 331:41
2. Briskin MJ, *et al.* 1993. *Nature* 363:461.
3. Berlin C, *et al.* 1993. *Cell* 74:185.
4. Lehnert K, *et al.* 1998. *Eur. J. Immunol.*