

PE anti-human/mouse integrin $\beta 7$

Catalog # / Size: 1205025 / 25 μ g
 1205030 / 100 μ g

Clone: FIB27

Isotype: Rat IgG2a, κ


Immunogen: TK1 cells

Reactivity: Human, Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 splenocytes stained with FIB27 PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

- Application References:**
1. Andrew DP, *et al.* 1994. *J. Immunol.* 153:3847. (Block)
 2. Berlin C, *et al.* 1993. *Cell* 74:185. (Block)
 3. Tidswell M, *et al.* 1997. *J. Immunol.* 159:1497. (Block)

Description: Integrin $\beta 7$ is a 130 kD glycoprotein, also known as integrin βp . It is a member of the Ig superfamily. In association with integrin $\alpha 4$ or αE chain, $\beta 7$ forms $\alpha 4/\beta 7$ or $\alpha E/\beta 7$ heterodimer. $\alpha 4/\beta 7$ (CD49d/ $\beta 7$, LPAM-1) is expressed on majority of peripheral lymphocytes, small subsets of thymocytes and bone marrow progenitors. LPAM-1 binds to several ligands, VCAM-1, MAdCAM-1 and fibronectin, and is involved in lymphocyte adhesion, some hematopoietic progenitor cells migration. $\alpha E/\beta 7$ (CD103/ $\beta 7$, $\alpha_{IEL}/\beta 7$) is expressed on intestinal intraepithelial lymphocytes (IEL), dendritic epidermal T cells, T regulatory cells, a subset of CD8+ T cells in lymph nodes and lamina propria. CD103/ $\beta 7$ complex is thought to play a role in lymphocyte retention via interaction with its lignd E-Cadherin. The FIB27 antibody has been reported to react with mouse and human $\beta 7$ integrin and to block $\beta 7$ integrin-mediated cell adhesion in *in vitro* and *in vivo* studies.

- Antigen References:**
1. Andrew DP, *et al.* 1994. *J. Immunol.* 153:3847.
 2. Picarella D, *et al.* 1997. *J. Immunol.* 158:2099.
 3. Lefrancois L, *et al.* 1994. *Eur. J. Immunol.* 24:635
 4. Cepek KL, *et al.*