


**PerCP/Cy5.5 anti-mouse CD43 Activation-Associated Glycoform**

<b>Catalog # / Size:</b>	1206120 / 100 µg 1206115 / 25 µg
<b>Clone:</b>	1B11
<b>Isotype:</b>	Rat IgG2a, κ
<b>Immunogen:</b>	Mouse WEHI 274.3 myeloid tumor cells
<b>Reactivity:</b>	Mouse
<b>Preparation:</b>	The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.2



Balb/c mouse splenocytes stained with 1B11 PerCP/Cy5.5

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 microg per 10<sup>6</sup> cells in 100 microL. It is recommended that the reagent be titrated for optimal performance for each application.

\* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>1,2</sup>, Western blotting<sup>1,2</sup>, and immunohistochemistry of acetone-fixed frozen sections.

**Application References:**

1. Jones AT, *et al.* 1994. *J. Immunol.* 153:3426. (IP, WB)
2. Carlow DA, *et al.* 1999. *J. Immunol.* 163:1441. (IP, WB)
3. Onami TM, *et al.* 2002. *J. Immunol.* 168:6022.
4. van der Most RG, *et al.* 2003. *Intl. Immunol.* 15:119.
5. Chu VT, *et al.* 2007. *J. Immunol.* 179:5947.
6. Lang A, *et al.* 2008. *J. Immunol.* 180:4848. [PubMed](#)
7. Gibbert K, *et al.* 2010. *J. Immunol.* 185:6179. [PubMed](#)

**Description:** CD43, also known as Ly-48, Leukosialin, Sialophorin, Leukocyte Sialoglycoprotein, and gp115, is a large single chain of type I transmembrane glycoprotein with abundant O-glycosylation and sialylation sites. Due to variable glycosylation and sialylation, two isoforms of CD43 have been identified. The 115 kD glycoform of CD43 is expressed on most hematopoietic cells including T lymphocytes, NK cells, monocytes, granulocytes, platelets, and CD5<sup>+</sup> B cells. It is not present on resting B cells and erythrocytes. While the 130 kD glycoform is thought to be activation-associated form primarily expressed on myeloid cells, pre-B cells, and activated T cells. It has been reported that CD43 binds to CD54 and Siglec-1. CD43 plays dual roles in cell adhesion and anti-adhesion, as well as costimulation of T cell

activation and survival, and induction of apoptosis of T cells and hematopoietic progenitors. The 1B11 antibody reacts with the activation-associated glycoform of CD43. The epitope recognized by 1B11 is also shared with desialylated CD45RB. This antibody is useful for differentiation of effector CD8 T cells and memory T cells.

**Antigen  
References:**

1. van den Berg TK, *et al.* 2001. *J. Immunol.* 166:3637.
2. Moore T, *et al.* 1994. *J. Immunol.* 153:4978.
3. Onami TM, *et al.* 2002. *J. Immunol.* 168:6022.
4. Tong J, *et al.* 2004.