PE anti-mouse CD309 (VEGFR2, Flk-1)

Catalog # / Size: 1209530 / 100 μg

1209525 / 25 μg

Clone: 89B3A5

Isotype: Rat IgG2a, κ

Immunogen: Rat-1 cells transfected with full-length

mouse Flk

Reactivity: Mouse

Preparation: The antibody was purified by affinity


chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



Mouse FLK-1 transfected cells stained with 89B3A5 PE

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is ≤ 1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application References:

1. Kaburn N, et al. 1997. Development 124:2039.

Description: The 89B3A5 antibody recognizes mouse CD309 also known as vascular

endothelial growth factor receptor 2, VEGFR2, KDR, protein tyrosine kinase receptor flk-1, and fetal liver kinase-1. Flk-1 is a member of the tyrosine protein kinase family, sub-family CSF-1/PDGF, that contains a single pass transmembrane receptor with a protein kinase domain and seven immunoglobulin-like domains in the extracellular region. Flk-1 is expressed at high levels in adult heart, lung, kidney, brain, and skeletal muscle; other tissues express at lower levels. Flk-1 is a receptor for VEGF or VEGFC; ligand binding plays a key role in vascular

development and vascular permeability. The 89B3A5 antibody has been shown to

be useful for flow cytometry.

Antigen

1. Patterson C, et al. 1995. J. Biol. Chem. 270:23111.

References: 2. Quinn TP, et al. 1993. Proc. Natl. Acad. Sci. USA 90:7533.