


Alexa Fluor® 488 anti-mouse Ly-6A/E (Sca-1)

Catalog # / Size: 1212580 / 100 µg
Clone: E13-161.7
Isotype: Rat IgG2a, κ
Immunogen: mouse pre-T cells
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 488 under optimal conditions.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



C57BL/6 splenocytes stained with E13-161.7 Alexa Fluor® 488

Applications:

Applications: Immunofluorescence
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.

Application Notes: Additional reported applications (for the relevant formats) include: complement-mediated cytotoxicity¹, immunohistochemical staining of frozen sections² and Bouin-fixed, paraffin-embedded samples⁴, immunofluorescence microscopy⁴, and immunoprecipitation³.

The two Sca-1 recognizing clones [D7](#) and E13-161.7 have been shown to bind distinct epitopes due to the inability of D7 to block the binding of E13-161.7.⁷

Application References:

1. Aihara Y, *et al.* 1986. *Eur. J. Immunol.* 16:1391. (CMCD)
2. Spangrude GJ, *et al.* 1988. *J. Immunol.* 141:3697. (IHC)
3. van de Rijn M, *et al.* 1989. *P. Natl. Acad. Sci. USA* 86:4634. (IP)
4. van Bragt MPA, *et al.* 2005. *Biol. Reprod.* 73:634. (IHC, IF)
5. Rosas M, *et al.* 2010. *J. Leukoc. Biol.* 88:169. [PubMed](#)
6. Felthaus O, *et al.* 2010. *Neurosci Lett.* 471:179. [PubMed](#)
7. Bamezai A and Rock KL. 1995. *Proc. Natl. Acad. Sci. USA* 92:4294.
8. Flaczyk A, *et al.* 2013. *J. Immunol.* 191:2503. [PubMed](#)

Description: Ly-6A/E is an 18 kD member of the Ly-6 multigene family also known as Sca-1. Ly6A/E is a glycosylphosphatidylinositol (GPI)-linked protein expressed on hematopoietic stem cells. In mice expressing the Ly-6.2 haplotype (e.g., AKR, C57BL, C57BR, DBA/2, SJL, SWR, and 129), Ly-6A/E is also expressed on peripheral B lymphocytes and thymic and peripheral T lymphocytes. Strains

expressing the Ly-6.1 haplotype (e.g., BALB/c, CBA, C3H/He, DBA/1, and NZB) have low Ly-6A/E expression on resting peripheral lymphocytes. The expression of Ly-6A/E on lymphocytes is upregulated upon activation from both Ly6.1 and Ly6.2 haplotype mice. Ly-6A/E is thought to be involved in the regulation of both T and B cell responses. The E13-161.7 antibody has been reported to be useful for immunohistochemical staining of frozen sections, immunoprecipitation, flow cytometry, and complement-mediated cytotoxicity.

**Antigen
References:**

1. Rock KL, *et al.* 1989. *Immunol. Rev.* 111:195.
2. Morrison SJ, *et al.* 1994. *Immunity* 1:661.
3. Spangrude GJ, *et al.* 1988. *J. Immunol.* 141:3697.
4. Malek T, *et al.* 1986. <