

**PE/Cy7 anti-mouse F4/80**

**Catalog # / Size:** 1215565 / 25 µg  
1215570 / 100 µg

**Clone:** BM8

**Isotype:** Rat IgG2a, κ


**Immunogen:** Murine macrophages

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



Thioglycolate-elicited BALB/c mouse peritoneal macrophages stained with BM8 PE/Cy7

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections<sup>1,2</sup> and formalin-fixed paraffin-embedded sections<sup>6,7</sup>, and Western blotting.

**Application References:**

- Schaller E, *et al.* 2002. *Mol. Cell. Biol.* 22:8035. (IHC)
- Stevceva L, *et al.* 2001. *BMC Clin Pathol.* 1:3. (IHC)
- Kobayashi M, *et al.* 2008. *J. Leukoc. Biol.* 83:1354. [PubMed](#)
- Poeckel D, *et al.* 2009. *J. Biol Chem.* 284:21077. [PubMed](#)
- Glass AM, *et al.* 2013. *J. Immunol.* 190:4830. [PubMed](#)
- Koehm S, *et al.* 2007. *J. Allergy Clin. Immunol.* 120:570. (IHC)
- Rankin AL, *et al.* 2010. *J. Immunol.* 184:1526. (IHC)
- Sasi SP, *et al.* 2014. *J Biol Chem.* 289:14178. [PubMed](#)
- Thakus VS, *et al.* 2014. *Toxicol Lett.* 230:322. [PubMed](#)
- Giordano C, *et al.* 2015. *Hum Mol Genet.* [PubMed](#)
- Hamilton JA, *et al.* 2015. *J Immunol.* 194:5022. [PubMed](#)
- Doni A, *et al.* 2015. *J Exp Med.* 212:905. [PubMed](#)
- Sharma SK, *et al.* 2015. *J Immunol.* 194:5529. [PubMed](#)

**Description:** F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on the majority of tissue macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node and Peyer's patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has

not been identified, but it has been reported that F4/80 is required for induction of CD8<sup>+</sup> T cells-mediated peripheral tolerance.

**Antigen  
References:**

1. Austy JM and Gordon S. 1981. *Eur. J. Immunol.* 11:805.
2. Hume DA, *et al.* 1983. *J. Exp. Med.* 158:1522.
3. Ruedl C, *et al.* 1996. *Eur. J. Immunol.* 26:1801.
4. McKnight AJ, *et al.* 1996