


PE/Dazzle™ 594 anti-mouse F4/80

Catalog # / Size:	1215725 / 25 µg 1215730 / 100 µg
Clone:	BM8
Isotype:	Rat IgG2a, κ
Immunogen:	Murine macrophages
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	0.2



Thioglycolate-elicited BALB/c mouse peritoneal macrophages were stained with F4/80 (clone BM8) PE/Dazzle™ 594 (filled histogram) or rat IgG2a, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application. * PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.
Application Notes:	Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections ^{1,2} and formalin-fixed paraffin-embedded sections ^{6,7} , and Western blotting.
Application References:	<ol style="list-style-type: none">Schaller E, <i>et al.</i> 2002. <i>Mol. Cell. Biol.</i> 22:8035. (IHC)Stevceva L, <i>et al.</i> 2001. <i>BMC Clin Pathol.</i> 1:3. (IHC)Kobayashi M, <i>et al.</i> 2008. <i>J. Leukoc. Biol.</i> 83:1354. PubMedPoeckel D, <i>et al.</i> 2009. <i>J. Biol Chem.</i> 284:21077. PubMedGlass AM, <i>et al.</i> 2013. <i>J. Immunol.</i> 190:4830. PubMedKoehm S, <i>et al.</i> 2007. <i>J. Allergy Clin. Immunol.</i> 120:570. (IHC)Rankin AL, <i>et al.</i> 2010. <i>J. Immunol.</i> 184:1526. (IHC)Sasi SP, <i>et al.</i> 2014. <i>J Biol Chem.</i> 289:14178. PubMedThakus VS, <i>et al.</i> 2014. <i>Toxicol Lett.</i> 230:322. PubMed

Description: F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on the majority of tissue macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node and Peyer's

patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been identified, but it has been reported that F4/80 is required for induction of CD8⁺ T cells-mediated peripheral tolerance.

**Antigen
References:**

1. Austy JM and Gordon S. 1981. *Eur. J. Immunol.* 11:805.
2. Hume DA, *et al.* 1983. *J. Exp. Med.* 158:1522.
3. Ruedl C, *et al.* 1996. *Eur. J. Immunol.* 26:1801.
4. McKnight AJ, *et al.* 1996