

APC/Fire™ 810 anti-mouse F4/80

Catalog # / Size: 1215825 / 25 µg
1215830 / 100 µg

Clone: BM8

Isotype: Rat IgG2a, κ


Immunogen: Murine macrophages

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 810 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Concentration: 0.2 mg/mL



Thioglycolate-elicited Balb/c mouse peritoneal macrophages were stained with anti-mouse CD11b FITC and anti-mouse F4/80 (clone BM8) APC/Fire™ 810 (left) or rat IgG2a, κ APC/Fire™ 810 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* APC/Fire™ 810 has a maximum excitation of 650 nm and a maximum emission of 810 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections^{1,2} and formalin-fixed paraffin-embedded sections^{6,7}, and Western blotting.

- Application References:**
1. Schaller E, *et al.* 2002. *Mol. Cell. Biol.* 22:8035. (IHC)
 2. Stevceva L, *et al.* 2001. *BMC Clin Pathol.* 1:3. (IHC)
 3. Kobayashi M, *et al.* 2008. *J. Leukoc. Biol.* 83:1354. [PubMed](#)
 4. Poeckel D, *et al.* 2009. *J. Biol Chem.* 284:21077. [PubMed](#)
 5. Glass AM, *et al.* 2013. *J. Immunol.* 190:4830. [PubMed](#)
 6. Koehm S, *et al.* 2007. *J. Allergy Clin. Immunol.* 120:570. (IHC)
 7. Rankin AL, *et al.* 2010. *J. Immunol.* 184:1526. (IHC)
 8. Sasi SP, *et al.* 2014. *J Biol Chem.* 289:14178. [PubMed](#)
 9. Thakus VS, *et al.* 2014. *Toxicol Lett.* 230:322. [PubMed](#)
 10. Watson NB, *et al.* 2015. *J Immunol.* 194:2796. [PubMed](#)
 11. Hirakawa H, *et al.* 2015. *PLoS One.* 10:119360. [PubMed](#)

Description: F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on the majority of tissue macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node and Peyer's patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been identified, but it has been reported that F4/80 is required for induction of CD8⁺ T cells-mediated peripheral tolerance.

Antigen
References:

1. Austy JM and Gordon S. 1981. *Eur. J. Immunol.* 11:805.
2. Hume DA, *et al.* 1983. *J. Exp. Med.* 158:1522.
3. Ruedl C, *et al.* 1996. *Eur. J. Immunol.* 26:1801.
4. McKnight AJ, *et al.* 1996. *J. Biol. Chem.* 271:486.
5. Lin HH, *et al.* 2005. *J. Exp. Med.* 201:1615.