

**PE/Cy7 anti-mouse CD14**

**Catalog # / Size:** 1216575 / 25 µg  
1216580 / 100 µg

**Clone:** Sa14-2

**Isotype:** Rat IgG2a, κ


**Immunogen:** Mouse thymus or spleen

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



Thioglycolate-elicited Balb/c mouse peritoneal macrophages stained with Sa14-2 PE/Cy7

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:** 1. Wallner S, *et al.* 2013. *PLoS One*. 8:65178. [PubMed](#)

**Description:** CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein also known as LPS receptor. CD14 is expressed on macrophages, dendritic cells, Kupffer cells, hepatocytes, and granulocytes. As a high-affinity receptor for LPS-LBP (LPS-binding protein) complex, CD14, in association with Toll-like Receptor 4 (TLR4) or 2 (TLR2), is involved in the clearance of gram-negative pathogens.

**Antigen References:**

1. Stocks S, *et al.* 1990. *Biochem. J.* 268:275.
2. Akashi S, *et al.* 2003. *J. Exp. Med.* 198:1035.
3. Matsuura K, *et al.* 1994. *J. Exp. Med.* 179:1671.
4. Liu S, *et al.* 1998.