## PE/Cy5 anti-mouse CD40

**Catalog # / Size:** 1223085 / 25 μg

1223090 / 100 µg

**Clone:** 3/23

**Isotype:** Rat IgG2a, κ

Immunogen: Recombinant mouse CD40 protein

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity


chromatography, and conjugated with PE/Cy5 under optimal conditions. The solution is free of unconjugated PE/Cy5

and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 splenocytes stained with

3/23 PE/Cy5

## **Applications:**

**Applications:** Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is  $\leq 0.25$  microg per  $10^6$  cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

**Application** 

Notes:

The LEAF  $^{\text{TM}}$  purified antibody (Endotoxin <0.1 EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for functional assays (Cat. No. 124604). For highly sensitive assays, we recommend Ultra-LEAF  $^{\text{TM}}$  purified antibody (Cat. No. 124628) with a lower endotoxin limit than standard LEAF  $^{\text{TM}}$  purified antibodies (Endotoxin <0.01 EU/ $\mu$ size  $\mu$ s)

EU/microg).

Application References:

Hasbold J, et al. 1994. Eur. J. Immunol. 24:1835.
Bourgeois C, et al. 2002. Science 297:2060.

**Description:** 

CD40 is a 48 kD type I transmembrane glycoprotein also known as Bp50. It is a member of the tumor necrosis factor receptor (TNFR) superfamily and is expressed on B cells, basal epithelial cells, macrophages, follicular dendritic cells, endothelial cells, and a subset of CD34<sup>+</sup> hematopoietic progenitors. CD40 regulates B cell development/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.

Antigen References: 1. Grewal IS, et al. 1998. Annu Rev Immunol 16:111.