## **Product Data Sheet**

## PE/Cy7 anti-mouse FR4 (Folate Receptor 4)

Catalog # / Size: 1225060 / 100 μg

Clone: 12A5

**Isotype:** Rat IgG2b, κ

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity


chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7

and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

**Concentration:** 0.2



C57BL/6 mouse splenocytes stained with CD3 (145-2C11) APC and 12A5 PE/CY7

## **Applications:**

**Applications:** Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is ≤ 0.25 microg per 106 cells in 100 microL volume. It is

recommended that the reagent be titrated for optimal performance for each

application.

Application

References:

1. Yamaguchi T et al. 1007: Immunity 27(1)145

**Description:** Folate receptor 4 (FR4) is a surface receptor for folic acid (Vitamin B9).It was

recently reported that FR4 is highly and constitutively expressed on mouse CD4<sup>+</sup> CD25<sup>+</sup> natural regulatory T cells (Treg). The high expression of this surface marker on Treg can be used in combination with CD4 and CD25 to distinguish Treg from other types of T cells. Since FR4 is a surface marker, this antibody may be used in combination with anti-CD4 and/or CD25 to sort living T reg cells for

functional studies.

Antigen

1. Yamaguchi T *et al* 2007. *Immunity* 27:145

References: 2. Spiegelstein O et al. 2000 Gene 258:117