Alexa Fluor® 700 anti-mouse CD8b

Catalog # / Size: 1233090 / 100 µg

1233085 / 25 µg

Clone: YTS156.7.7 Isotype: Rat IgG2b, ĸ

Immunogen: Mouse thymocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity


> chromatography and conjugated with Alexa Fluor® 700 under optimal

conditions.

Phosphate-buffered solution, pH 7.2, Formulation:

containing 0.09% sodium azide.

Concentration: 0.5



C57BL/6 mouse splenocyte stained with CD8b (clone YTS156.7.7) Alexa Fluor® 700 (filled histogram) or rat IgG2a, κ Alexa Fluor® 700 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes:

Additional reported applications (for the relevant formats) include: immunohistochemistry of acetone-fixed frozen tissue sections.

Application References:

1. McNical AM, et al. 2007. Eur. J. Immunol. 37:1634.

Description: CD8b is the 32 kD β chain of CD8, also known as Lyt-3 or Ly-3. It is a member of

the Ig superfamily expressed as a heterodimer with the CD8 α chain on a subset of MHC class I-restricted T cells and most thymocytes. CD8 is a co-receptor for the

TCR complex involved in T cell activation.

Antigen References: 1. Barclay A, et al. 1997. The Leukocyte antigen Facts Book Academic Press.

2. Zamovska R. 1994. Immunity 1:243-246.

3. Ellmeier W, et al. 1999. Annu. Rev Immunol 17:523.

4. Ledbetter IA, et al.