

**FITC anti-mouse Podoplanin**

**Catalog # / Size:** 1237075 / 25 µg  
1237080 / 100 µg

**Clone:** 8.1.1


**Isotype:** Hamster IgG

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** Lot-specific



Mouse thymic epithelial stromal cell line TE-71 stained with Podoplanin (clone 8.1.1) FITC (filled histogram) or left unstained (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunohistochemistry<sup>6</sup>.

- Application References:**
1. Farr A, *et al.* 1992. *J. Histochem. Cytochem.* 40:651.
  2. Farr AG, *et al.* 1992. *J. Exp. Med.* 176:1477.
  3. Bekiaris V, *et al.* 2008. *J. Immunol.* 180:6768.
  4. Algars A, *et al.* 2011. *Blood* 117:4387. [PubMed](#)
  5. Reis VO, *et al.* 2012. *Immunobiology.* 217:831. [PubMed](#)
  6. Kaji C, *et al.* 2012. *Acta. Histochem. Cytochem.* 45:227. (IHC)
  7. Kretschmer S, *et al.* 2013. *PLoS One.* 8:e52201. [PubMed](#).

**Description:** The mucin-type glycoprotein podoplanin is thought to be involved in the development of the lymphatic vascular system. Podoplanin is named after its expression in the kidney glomerular epithelial cells (podocytes). It has a potential role in tumor progression.

- Antigen References:**
1. Farr A, *et al.* 1992. *J. Histochem. Cytochem.* 40:651.
  2. Schacht V, *et al.* 2005. *Am. J. Pathol.* 166:913.

