## **Product Data Sheet**

## APC/Cy7 anti-mouse Ly-6C

Catalog # / Size: 1240130 / 100 μg

1240125 / 25 µg

Clone: HK1.4

**Isotype:** Rat IgG2c, κ

Immunogen: L3 cloned CTL cells

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity


chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7

and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

**Concentration:** 0.2



C57BL/6 bone marrow cells stained with HK1.4 APC/Cy7

## **Applications:**

**Applications:** Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25$  microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes:

Clone HK1.4 does not block the binding of clone RB6-8C5<sup>8</sup>.

Additional reported applications (for relevant formats of this clone) include: *in vitro* activation of T cells<sup>1-3</sup> and immunohistochemistry of frozen sections4.

Application References:

- 1. Jutila MA, et al. 1988. Eur. J. Immunol. 18:1819. (Activ)
- 2. Herold KC, et al. 1990. Diabetes 39:815. (Activ)
- 3. Havran WL, et al. 1988. J. Immunol. 140:1034 (Activ)
- 4. Flanagan K, et al. 2008. J. Immunol. 180:3874. (IHC)
- 5. Makaroff LE, et al. 2009. P. Natl. Acad. Sci. USA 106:4799. (FC)
- 6. Zuber J, *et al.* 2009. *Genes Dev.* 23:877. (FC) <u>PubMed</u>
- 7. Ribechini E, et al. 2009. Eur. J. Immunol. 39:3538.
- 8. Ma C, et al. 2012. J. Leukoc. Biol. 92:1199.
- 9. Watson NB, et al. 2015. J Immunol. 194:2796. PubMed

**Description:** Most hematopoietic cells express one or more members of Ly-6 family. The

expression of Ly-6 varies with development stage and activation. Ly-6C is a 14-17 kD GPI-linked surface protein expressed on mouse monocyte/macrophage cells, endothelial cells, neutrophils, and some T cell subsets. Ly-6C is reported to be an

indicator of memory CD8 $^{+}$  T cells.

Antigen

1. Jutila MA, et al. 1988. Eur. J. Immunol. 18:1819.

References: 2

2. Cerwenka A, et al. 1998. J. Immunol. 161:97.