

**PE/Dazzle™ 594 anti-mouse Ly-6C**

**Catalog # / Size:** 1240220 / 100 µg  
1240215 / 25 µg

**Clone:** HK1.4

**Isotype:** Rat IgG2c, κ


**Immunogen:** L3 cloned CTL cells

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



C57BL/6 mouse bone marrow cells were stained with Ly-6C (clone HK1.4) PE/Dazzle™ 594 (filled histogram) or rat IgG2a, κ PE/Dazzle™ 594 isotype control (open histogram). Data shown was gated on the myeloid population.

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.06 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

**Application Notes:** Clone HK1.4 does not block the binding of clone RB6-8C5<sup>8</sup>.  
Additional reported applications (for relevant formats of this clone) include: *in vitro* activation of T cells<sup>1-3</sup> and immunohistochemistry of frozen sections<sup>4</sup>.

**Application References:**

1. Jutila MA, *et al.* 1988. *Eur. J. Immunol.* 18:1819. (Activ)
2. Herold KC, *et al.* 1990. *Diabetes* 39:815. (Activ)
3. Havran WL, *et al.* 1988. *J. Immunol.* 140:1034 (Activ)
4. Flanagan K, *et al.* 2008. *J. Immunol.* 180:3874. (IHC)
5. Makaroff LE, *et al.* 2009. *P. Natl. Acad. Sci. USA* 106:4799. (FC)
6. Zuber J, *et al.* 2009. *Genes Dev.* 23:877. (FC) [PubMed](#)
7. Ribechini E, *et al.* 2009. *Eur. J. Immunol.* 39:3538.
8. Ma C, *et al.* 2012. *J. Leukoc. Biol.* 92:1199.
9. Watson NB, *et al.* 2015. *J Immunol.* 194:2796. [PubMed](#)

**Description:** Most hematopoietic cells express one or more members of Ly-6 family. The expression of Ly-6 varies with development stage and activation. Ly-6C is a 14-17 kD GPI-linked surface protein expressed on mouse monocyte/macrophage cells, endothelial cells, neutrophils, and some T cell subsets. Ly-6C is reported to be an

indicator of memory CD8<sup>+</sup> T cells.

- Antigen** 1. Jutila MA, *et al.* 1988. *Eur. J. Immunol.* 18:1819.
- References:** 2. Cerwenka A, *et al.* 1998. *J. Immunol.* 161:97.