Product Data Sheet

PE anti-mouse CD226 (DNAM-1)

Catalog # / Size:	1244030 / 100 μg 1244025 / 25 μg	2
Clone:	10E5	
Isotype:	Rat IgG2b, к	3
Immunogen:	Th1 polarized T cell clones	Ω ™
Reactivity:	Mouse	
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.	CD8 (53-6.7) FITC C57BL/6 splenocytes stained with CD8 (53-6.7) FITC and 10E5 PE
Concentration:	0.2	

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application References:	1. Dardalhon V, <i>et al.</i> 2005. <i>J. Immunol.</i> 175:1558 2. Huang Z, <i>et al.</i> 2011. <i>J Immunol.</i> 186:3421. <u>PubMed</u>
Description:	CD226 (DNAM-1) is constitutively expressed on native CD8 ⁺ cells and on CD4 ⁺ T cells, macrophages and NK cells.This antibody (10E5) was reported to bind about 40% of inactivated CD4 ⁺ cells and binds only to differentiated Th1 cells, but not to Th2 or Th0 cells.It is also reported to suppress antigen-specific T cell expansion and EAE (experimental allergic encephalitis) mediated by Th1 cells.
Antigen References:	1.Tahara Hanaoka S, <i>et al.</i> 2006. <i>Blood</i> 107:1491 2. Shibuya K, <i>et al.</i> 2003. <i>J. Exp. Med.</i> 198:1829