

**APC anti-mouse CD196 (CCR6)**

**Catalog # / Size:** 1249070 / 100 µg  
1249065 / 25 µg

**Clone:** 29-2L17

**Isotype:** Hamster IgG


**Immunogen:** N-terminal peptide (aa. 2-38) of mouse CCR6 fused with GST

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2




C57BL/6 splenocytes stained with 29-2L17 APC and CD45R/B220 (RA3-6B2) PE

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



C57BL/6 splenocytes stained with Armenian Hamster IgG isotype control APC and CD45R/B220 (RA3-6B2) PE

**Application References:** 1. Mony JT, *et al.* 2014. *Mult Scler.* [PubMed](#)

**Description:** CCR6 is a G-protein linked chemokine receptor which binds the chemokine CCL20/MIP-3a. It is expressed on B lymphocytes and certain subsets of dendritic cells and T cells. CCR6 is reported to be involved in mucosal immune response and lymphocyte migration and homeostasis.

**Antigen References:** 1. Rossi D, *et al.* 1997. *J. Immunol.* 158:1033.  
2. Lukacs N, *et al.* 2001. *J. Exp. Med.* 194:551.  
3. Cook DN, *et al.* 2000. *Immunity* 12:495