

Brilliant Violet 421™ anti-mouse CD196 (CCR6)

Catalog # / Size: 1249090 / 500 µl
1249085 / 125 µl

Clone: 29-2L17

Isotype: Hamster IgG


Immunogen: N-terminal peptide (aa. 2-38) of mouse CCR6 fused with GST

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific




C57BL/6 mouse splenocytes were stained with CD45R/B220 FITC and CCR6 (clone 29-2L17) Brilliant Violet 421™ (top) or Armenian hamster IgG Brilliant Violet 421™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Description: CCR6 is a G-protein linked chemokine receptor which binds the chemokine CCL20/MIP-3a. It is expressed on B lymphocytes and certain subsets of dendritic cells and T cells. CCR6 is reported to be involved in mucosal immune response and lymphocyte migration and homeostasis.

Antigen
References:

1. Rossi D, *et al.* 1997. *J. Immunol.* 158:1033.
2. Lukacs N, *et al.* 2001. *J. Exp. Med.* 194:551.
3. Cook DN, *et al.* 2000. *Immunity* 12:495