

PerCP/Cy5.5 anti-mouse CD79b (Igβ)

Catalog # / Size: 1264045 / 25 µg
1264050 / 100 µg

Clone: HM79-12


Isotype: Hamster IgG

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: Lot-specific



C57BL/6 bone marrow cells stained with CD45R/B220 PE and CD79b (clone HM79-12) PerCP/Cy5.5 (top) or Armenian Hamster IgG isotype control PerCP/Cy5.5 (bottom).


Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.75 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.



- Application References:**
1. Gong S, *et al.* 1996. *Science*. 272:411.
 2. Nagata K, *et al.* 1997. *Immunity*. 7:559.
 3. Papavasiliou F, *et al.* 1995. *Science*. 268:408.

Description: Mouse CD79b (Ig β chain) is a 35-40kD transmembrane protein that forms a heterodimer with CD79a (30-35 kD, Ig α chain). The CD79b and CD79a heterodimers are associated with surface IgM to form the B-cell receptor (BCR) that is necessary for signal transduction via the BCR in mature B cells. CD79b participates in the signal transduction involved in development of B cells as well. It was reported that association between CD79b/CD79a with IgM is essential in inducing both the transition from progenitor to precursor B cells and subsequent allelic exclusion. Ig β knockout mice had a complete block in B cell development at the immature CD43⁺B220⁺ stage. The HM79b-12 clone reacts with an extracellular epitope of CD79b or Ig β .