

**Purified anti-mouse CD41**

**Catalog # / Size:** 1269505 / 50 µg  
1269510 / 500 µg

**Clone:** MWRReg30

**Isotype:** Rat IgG1, κ


**Immunogen:** Mouse platelets

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5



C57 BL/6 platelets were stained with CD41 (clone:MWRReg30) (filled histogram) or rat IgG1, κ isotype control (open histogram), followed by anti-rat IgG FITC.

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: depletion of platelets and functional assay *in vivo*.<sup>4,7</sup> The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for *in vivo* studies (Cat. No. 133910).

- Application References:**
1. Nieswandt B, *et al.* 1999. *Blood* 94:684.
  2. Teeling JL, *et al.* 2001. *Blood* 98:1095.
  3. Bertrand JY, *et al.* 2005. *P. Natl. Acad. Sci. USA* 102:134.
  4. Nocito A, *et al.* 2007. *Hepatology* 45:369. (Deplete)
  5. Sullivan BP, *et al.* 2010. *Toxicol. Sci.* 115:286. (Deplete) [PubMed](#)
  6. van der Heyde HC, *et al.* 2005. *Blood* 105:1956. (FA)
  7. Marjon KD, *et al.* 2009. *J. Immunol.* 182:1397. (Deplete)

**Description:** CD41, also known as integrin α2b and GPIIb, is a transmembrane glycoprotein that is expressed by platelets and megakaryocytes. It was reported that CD41 is also expressed on hematopoietic progenitors. CD41 associates with CD61 (integrin β3) to form complexes that interact with fibrinogen, fibronectin, von Willebrand factor, and thrombin. CD41 is required for platelet adhesion and aggregation. Defect of CD41 leads to disorders of coagulation.

- Antigen References:**
1. Bakewell SJ, *et al.* 2003. *P. Natl. Acad. Sci. USA* 100:14205.
  2. Phillips DR, *et al.* 1991. *Cell.* 65:359.