## SONY

## **Product Data Sheet**

## Purified anti-mouse CD366 (Tim-3)

Catalog # / Size: 1270010 / 500 μg

Clone: B8.2C12
Isotype: Rat IgG1, κ

Immunogen: mTim-3 protein/Freund adjuvant

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity

chromatography.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

**Concentration:** 0.5

## **Applications:**

**Applications:** Flow Cytometry

**Recommended**Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application Notes:

Clone B8.2C12 only binds to the BALB/c allele of Tim-3.

Application

References:

1. del Rio ML, et al. 2011. Transpl. Int. 24:501. (FC) PubMed

**Description:** CD366 (Tim-3) is a transmembrane protein also known as T cell immunoglobulin

and mucin domain containing protein-3. Tim-3 is expressed at high levels on Th1 lymphocytes and CD11b<sup>+</sup> macrophages. Tim-3 has also been shown to exist as a soluble protein. Cells expressing Tim-3 are present at high levels in the CNS of animals at the onset of experimental autoimmune encephalomyelitis (EAE), a disease mediated by lymphocytes secreting Th1-like cytokines. Tim-3 has been proposed to inhibit Th1-mediated immune responses and promote immunological

tolerance.

Antigen References:

1. Sabatos CA, et al. 2003. Nat. Immunol. 4:1102

2. Kuchroo VK, et al. 2003. Nat. Rev. Immunol. 3:454

3. Mooney L, et al. 2002. Nature. 415:536

4. Rodriguez-Manzanet R, et al.