

Pacific Blue™ anti-mouse FcεRIα

Catalog # / Size: 1271565 / 25 µg
1271570 / 100 µg

Clone: MAR-1


Isotype: Hamster IgG

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with Pacific Blue™ under optimal conditions. The solution is free of unconjugated Pacific Blue™.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



Mouse mast cell line MC/9 stained with anti-mouse FcεRIα (clone MAR-1) Pacific Blue™

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume or 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: Additional reported applications (for relevant formats of this clone) include: depletion2, immunohistochemistry of frozen sections (OCT embedded2).

- Application References:**
1. Obata K, *et al.* 2007. *Blood* 110:913 (FC)
 2. Sokol CL, *et al.* 2008. *Nat. Immunol.* 9:310 (FC, Deplete, IHC)
 3. Chen J, *et al.* 2009. *J. Biol. Chem.* 284:5763 (FC)

Description: FcεRIα is a transmembrane protein belonging to the Ig superfamily. FcεRIα forms a tetrameric complex with one β and two γ-subunits. The FcεRI complex plays an important role in triggering IgE-mediated allergic reactions. It is abundantly expressed on mast and basophils and up-regulated by the presence of IgE. Following stimulation via FcεRIα, mast cells and basophils release bioactive chemical mediators such as histamine, resulting in the initiation of allergic reactions. Cross linking of the high-affinity receptor for IgE on tissue mast cells triggers immediate hypersensitivity with local symptoms. The MAR-1 monoclonal antibody reacts with the FcεRIα subunit.

- Antigen References:**
1. Arinobu Y, *et al.* 2005. *P. Natl. Acad. Sci. USA* 102:18105.
 2. Yamaguchi M, *et al.* 2001. *Int. Immunol.* 13:843.