
Product Data Sheet

PE anti-mouse CD135


Catalog # / Size:	1276525 / 50 μg 1276530 / 200 μg
Clone:	A2F10
Isotype:	Rat IgG2a, κ
Immunogen:	Mouse Flt3 transfected cell line
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	0.2



Bone marrow cells from C57BL/6 mouse stained with A2F10 PE and CD45R/B220 (RA3-6B2) FITC

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 microg per 10^6 cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



Bone marrow cells from C57BL/6 mouse stained with rat IgG2a, k PE and CD45R/B220 (RA3-6B2) FITC

Application	1. Sergejeva S, <i>et al.</i> 2004. <i>Blood</i> 103:1270.
References:	2. Auffray C, <i>et al.</i> 2009. <i>J. Exp. Med.</i> 206:595. 3. Chiba H, <i>et al.</i> 2013. <i>Am J Physiol Cell Physiol.</i> 305:693. <u>PubMed</u>

Description:	CD135, also known as Flk-2, Flt3, and Ly-72, is a type III tyrosine kinase receptor. It is expressed on early B lymphoid lineage cells in bone marrow, on primitive myeloid progenitors within the BM CD34+ cell population. Ligation of Flk-2 with Flt3 ligand regulates the growth of hematopoietic stem cells and promotes the survival of primitive hematopoietic progenitor cells with myeloid as well as B lymphoid potential. It was reported that the receptor tyrosine kinase Flt3 is required for dendritic cell development. Combined signaling through interleukin-7 receptors and Flt3 selectively promotes B-cell commitment and differentiation from uncommitted murine bone marrow progenitor cells.
	required for dendritic cell development. Combined signaling through interleukin-7 receptors and Flt3 selectively promotes B-cell commitment and differentiation

Antigen	1. Waskow C, <i>et al.Nat. Immunol.</i> 9:676
References:	2. Veiby OP, <i>et al.</i> 1996. <i>Blood</i> 88(4):1256
	3. Veiby OP, et al. 1996. J. Immunol. 157(7):2953
	4. Mattews W, et al. 1991. Cell. 65(7):114

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com