



Brilliant Violet 605™ anti-mouse CD115 (CSF-1R)

Catalog # / Size: 1277585 / 50 µg
Clone: AFS98
Isotype: Rat IgG2a, κ
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 605™ and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Concentration: 0.2



Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.

Thioglycolate-elicited C57BL/6 mouse peritoneal macrophages were stained with F4/80 APC and CD115 (clone AFS98) Brilliant Violet 605™ (top) or rat IgG2a, κ Brilliant Violet 605™ isotype control (bottom).

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by

U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: blocking of ligand binding¹. The LEAF™ purified antibody (Endotoxin <0.1 EU/microg, Azide-Free, 0.2 µm filtered) is recommended for functional assays.

Application References: 1. Sudo T, *et al.* 1995. *Oncogene* 11:2469.
2. Murayama T, *et al.* 1999. *Circulation* 99:1740.
3. Jaeger BN, *et al.* 2012. *J. Exp. Med.* 209:565. [PubMed](#)

Description: CSF-1R, also known as CD115 and M-CSFR, is a single-pass type I membrane protein and member of the platelet-derived growth factor receptor family. This c-fms (Fms proto-oncogene) gene product's natural ligands include M-CSF and IL-34. Structural studies of CD115 have described an Ig-like extracellular domain, a transmembrane domain, an intracellular juxtamembrane domain, a split tyrosine kinase domain, and a C-terminal tail receptor. Receptor activation induces homodimerization in addition to phosphorylation and ubiquitination of intracellular residues. CD115 directly influences tissue macrophage and osteoclast differentiation and proliferation. It is expressed on monocytes/macrophages, peritoneal exudate cells, plasmacytoid and conventional dendritic cells, and osteoclasts.

Antigen References: 1. Sudo T, *et al.* 1995 *Oncogene* 11:2469.
2. Murayama T, *et al.* 1999 *Circulation* 99:1740.
3. Goswami S, *et al.* 2005 *Cancer Res.* 65:5278.
4. Yu W, *et al.* 2008 *J. Leuko. Bio*