

**PE/Cy7 anti-mouse CD115 (CSF-1R)**

**Catalog # / Size:** 1277615 / 25 µg  
1277620 / 100 µg

**Clone:** AFS98


**Isotype:** Rat IgG2a, κ

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



Thioglycolate-elicited BALB/c peritoneal macrophages were stained with F4/80 FITC and CD115 (clone AFS98) PE/Cy7 (top) or rat IgG2a, κ PE/Cy7 isotype control (bottom).


**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.125 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** Additional reported applications (for the relevant formats) include: blocking of ligand binding<sup>1</sup>. The LEAF™ purified antibody (Endotoxin <0.1 EU/microg, Azide-Free, 0.2 µm filtered) is recommended for functional assays.



- Application** 1. Sudo T, *et al.* 1995. *Oncogene* 11:2469.  
**References:** 2. Murayama T, *et al.* 1999. *Circulation* 99:1740.  
3. Jaeger BN, *et al.* 2012. *J. Exp. Med.* 209:565. [PubMed](#)
- 

**Description:** CSF-1R, also known as CD115 and M-CSFR, is a single-pass type I membrane protein and member of the platelet-derived growth factor receptor family. This c-fms (Fms proto-oncogene) gene product's natural ligands include M-CSF and IL-34. Structural studies of CD115 have described an Ig-like extracellular domain, a transmembrane domain, an intracellular juxtamembrane domain, a split tyrosine kinase domain, and a C-terminal tail receptor. Receptor activation induces homodimerization in addition to phosphorylation and ubiquitination of intracellular residues. CD115 directly influences tissue macrophage and osteoclast differentiation and proliferation. It is expressed on monocytes/macrophages, peritoneal exudate cells, plasmacytoid and conventional dendritic cells, and osteoclasts.

- Antigen** 1. Sudo T, *et al.* 1995 *Oncogene* 11:2469.  
**References:** 2. Murayama T, *et al.* 1999 *Circulation* 99:1740.  
3. Goswami S, *et al.* 2005 *Cancer Res.* 65:5278.  
4. Yu W, *et al.* 2008 *J. Leuko. Bio*