
PE/Cy7 anti-mouse CD140a

Catalog # / Size: 1279560 / 100 µg
1279555 / 25 µg

Clone: APA5

Isotype: Rat IgG2a, κ

Reactivity: Mouse

Concentration: 0.2

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported (for relevant formats) applications include: Western Blot, blocking function², and immunohistochemical staining of paraffin and frozen sections. The LEAF™ purified antibody is recommended for functional assays.

Description: Platelet-derived growth factor receptor-α (PDGFR-α), CD140a, is one of two receptors for platelet-derived growth factors (PDGFs) and binds to all isoforms of PDGFs: PDGF-AA, PDGF-AB, and PDGF-BB. PDGFRα is a receptor tyrosine kinase that forms homodimers or heterodimers on the surface upon ligand binding and phosphorylates substrates. PDGFRs consist of either homodimers of α/α, β/β, or heterodimers of α/β. PDGF receptors, α and β, are single glycoproteins with intracellular tyrosine kinase domain. Their ligand, PDGF, is a mitogen for connective tissue and glial cells. CD140a is expressed on embryonic tissues and mesenchymal-derived cells of adult mice. PDGF plays a role in wound healing and acts as a chemoattractant for fibroblasts, smooth muscle cells, glial cells, monocytes, and neutrophils.