

**PerCP/Cy5.5 anti-mouse CD68**

**Catalog # / Size:** 1285045 / 25 µg  
1285050 / 100 µg

**Clone:** FA-11

**Isotype:** Rat IgG2a


**Immunogen:** Purified Con A receptor glycoproteins from the P815 cell line

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



Thioglycolate-elicited Balb/c peritoneal macrophages intracellularly stained with FA-11 PerCP/Cy5.5

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

**Application Notes:** Additional reported (for relevant formats) applications include: immunoprecipitation<sup>1, 2</sup>, Western Blot<sup>1, 2</sup>, and immunohistochemical staining of frozen section<sup>2</sup>.

**Application References:**

1. Silva RP, *et al.* 1999. *Biochem. J.* 338:687. (IP WB)
2. Rabinowitz SS, *et al.* 1991. *J. Exp. Med.* 174:827. (IP WB IHC)
3. Olver JA, *et al.* 2012. *Am J Physiol Renal Physiol.* 302:1362. [PubMed](#)
4. Gruber RC, *et al.* 2014. *J Neurosci.* 34:16320. [PubMed](#)

**Description:** Mouse CD68, also known as macrosialin, is an 85-115 kD member of the lysosomal-associated membrane protein (LAMP) family. It is a heavily glycosylated and predominantly intracellular protein, mainly in late endosomes. Macrosialin is the murine homolog to the human macrophage glycoprotein CD68. It is expressed on tissue macrophages, Langerhans cells and at low levels on dendritic cells. Lamp proteins may have functions relating to cell-cell interaction or cell-ligand interaction. The biological function of CD68 is not completely understood.

**Antigen References:**

1. Ramprasad MP, *et al.* 1996. *Proc. Natl. Acad. Sci. USA* 93:14833.
2. Smith MJ, *et al.* 1987. *J. Cell. Sci.* 87:113.