

PE anti-mouse CD335 (NKp46)

Catalog # / Size: 1288020 / 100 µg
1288015 / 25 µg

Clone: 29A1.4

Isotype: Rat IgG2a, κ


Immunogen: NKp46-IgG1 Fc fusion protein

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse splenocytes stained with 29A1.4 PE and NK1.1 (PK136) APC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of frozen tissue sections^{1,2} and *in vitro* activation of NK cells¹.

Application References:

1. Walzer T, *et al.* 2007. *P. Natl. Acad. Sci. USA* 104:3384. (FC, Activ)
2. Walzer T, *et al.* 2007. *Nat. Immunol.* 8:1337. (FC, Activ)
3. Guerriero JL, *et al.* 2011. *J. Immunol.* 186:3517. (IHC) [PubMed](#)

Description: CD335, also known as NKp46, is a single-pass type I membrane protein of 46 kD. It belongs to the natural cytotoxicity receptor (NCR) family and contains two Ig-like (immunoglobulin-like) domains. It's expression is restricted to NK cells and a subset of NKT cells; it's not expressed in CD1d-restricted NKT cells. CD335 is a receptor for viral hemagglutinins and heparan sulfate proteoglycans and is involved in NK cell activation.

Antigen References:

1. Colucci F and Cilio CM. 2010. *Nat. Immunol.* 125:60.
2. Caligiuri MA. 2008. *Blood* 112:461.
3. Colonna M. 2009. *Immunity* 31:15.