Product Data Sheet

PE/Dazzle™ 594 anti-mouse CD335 (NKp46)

Catalog # / Size: 1288145 / 25 μg

1288150 / 100 µg

Clone: 29A1.4

Isotype: Rat IgG2a, κ

Immunogen: NKP46-IgG1 Fc fusion protein

Reactivity: Mouse

Preparation: The antibody was purified by affinity


chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: Lot-specific



C57BL/6 splenocytes were stained with NK1.1 FITC and CD335 (clone 29A1.4) PE/Dazzle™ 594 (top) or rat IgG2a, κ PE/Dazzle™ 594 isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage:

Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.75 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum

emission of 610 nm.

Application Notes:

Additional reported applications (for the

relevant formats) include:

immunohistochemical staining of frozen tissue sections^{1,2} and *in vitro* activation of NK cells1.

of MK cells1.

Application References:


1. Walzer T, et al. 2007. P. Natl. Acad. Sci. USA 104:3384. (FC, Activ)

2. Walzer T, et al. 2007. Nat. Immunol. 8:1337. (FC, Activ)

3. Guerriero JL, et al. 2011. J. Immunol. 186:3517. (IHC) PubMed

Description:

CD335, also known as NKp46, is a single-pass type I membrane protein of 46 kD. It belongs to the natural cytotoxicity receptor (NCR) family and contains two Iglike (immunoglobulin-like) domains. It's expression is restricted to NK cells and a subset of NKT cells; it's not expressed in CD1d-restricted NKT cells. CD335 is a receptor for viral hemagglutinins and heparan sulfate proteoglycans and is involved in NK cell activation.



Antigen References: Colucci F and Cilio CM. 2010. *Nat. Immunol.* 125:60.
Caligiuri MA. 2008. *Blood* 112:461.

3. Colonna M. 2009. *Immunity* 31:15.