

PE/Cy7 anti-mouse CD26 (DPP-4)

Catalog # / Size: 1289045 / 25 µg
1289050 / 100 µg

Clone: H194-112

Isotype: Rat IgG2a, κ


Immunogen: BALB/c thymocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 splenocytes were stained with CD26 (clone H194-112) PE/Cy7 (filled histogram) or rat IgG2a, κ PE/Cy7 (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications include: *in vitro* activation of thymocytes¹, immunoprecipitation from cell lysates and cell-free supernatants^{1,2} and immunohistochemical staining of frozen tissue sections¹.

Application References:

1. Naquet P, *et al.* 1988. *J. Immunol.* 141:4101.
2. Vivier I, *et al.* 1991. *J. Immunol.* 147:447.
3. Sen A, *et al.* 2012. *PNAS* [PubMed](#)
3. Coleman CM, *et al.* 2014. *J Gen Virol.* 95:408. [PubMed](#)

Description: CD26, also known as DPP IV or THAM, is a 220 kD type II transmembrane homodimer. It consists of an α/β hydrolase domain and an eight-blade β-propeller domain. After proteolysis of the membrane-bound CD26, a soluble form of DPP IV is released. CD26 is expressed on thymocytes (development dependent), T cells, B cells, NK cells, and macrophages. It is involved in T cell costimulation, endothelial cell migration and proteolysis processes.

Antigen References:

1. Cooper KG, *et al.* 2009. *Infect. Immun.* 77:2447.
2. Eltzschig HK, *et al.* 2006. *Blood* 108:1602.

3. Peranteau WH, *et al.* 2006. *Blood* 108:4268.