


Brilliant Violet 421™ anti-mouse CD144 (VE-cadherin)**Catalog # / Size:** 1290065 / 50 µg**Clone:** BV13**Isotype:** Rat IgG1, κ**Immunogen:** VE-cadherin-Ig fusion protein**Reactivity:** Mouse**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).**Concentration:** 0.2

Mouse endothelial cells bEnd.3 were stained with CD144 (clone BV13) Brilliant Violet 421™ (filled histogram) or rat IgG1 Brilliant Violet 421™ isotype control (open histogram).

Applications:**Applications:** Flow Cytometry**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume or 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Clone BV13 recognizes an epitope between aa 45 and 56, and has a binding affinity of 5-15 nM.5 Additional reported applications (for relevant formats) include: Western blotting1, blocking of cell interactions *in vivo*1, and immunofluorescence microscopy4.**Application References:**

1. Corada M, *et al.* 1999. *P. Natl. Acad. Sci. USA* 96:9815. (WB, Block)
2. Liao F, *et al.* 2000. *Cancer Res.* 60:6805. (FC)
3. Crosby CV, *et al.* 2005. *Blood* 105:2771. (FC)
4. Liao F, *et al.* 2002. *Cancer Res.* 62:2567. (IF)
5. May C, *et al.* 2005. *Blood* 105:4337. (epitope)

Description: CD144, also known as vascular endothelial-cadherin (VE-cadherin), is a 120 kD

member of the type II Cadherin family. It is an endothelial specific hemophilic adhesion molecule involved in endothelial cell survival, migration, contact-dependent growth inhibition, and homophilic adhesion. VE-cadherin is essential for maintaining the integrity of the endothelial barrier *in vivo*.

**Antigen
References:**

1. Allport JR, *et al.* 2002. *J. Leukocyte Biol.* 71:821.
2. Hirashima M, *et al.* 2009. *Blood* 93:1253.
3. Matsuyoshi N, *et al.* 1997. *Proc. Assoc. Am. Physicians* 109:362.
4. Matsumura K