

PE anti-mouse CD205 (DEC-205)

Catalog # / Size: 1291065 / 25 µg
1291070 / 100 µg

Clone: NLDC-145

Isotype: Rat IgG2a, κ


Immunogen: Mouse lymph node tissue

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: NULL




C57BL/6 mouse bone marrow cells were stained with Ly-6G/Ly-6C APC and CD205 (clone NLDC-145) PE (top) or rat IgG2a PE isotype control (bottom). Data shown was gated on the myeloid population.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.03 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections¹, Western Blot¹⁻³, and immunoprecipitation of bone marrow dendritic cell extracts².



Application References:

1. Witmer-Pack MD, *et al.* 1995. *Cell. Immun.* 163:157. (IHC, WB)
2. Swiggard WJ, *et al.* 1995. *Cell. Immun.* 165:302. (WB, IP)
3. Bonifaz LC, *et al.* 2004. *J. Exp. Med.* 199:815. (WB)
4. Yamazaki S, *et al.* 2002. *J. Immunol.* 181:6923. (FC)
5. Bankoti J, *et al.* 2010. *Toxicol. Sci.* 115:422. (FC)

Description: CD205, also known as DEC-205, is a 205 kD integral membrane protein homologous to the macrophage mannose receptor. It is a type I cell surface protein that belong to the C-type lectin family. CD205 is expressed at high levels by dendritic cells and thymic epithelial cells. It is also expressed by a number of other cell types, such as B lymphocytes, macrophages, Langerhans cells, bone marrow stromal cells, granulocytes, epithelial cells of pulmonary airways, and the capillaries of the brain. CD205 is a novel endocytic receptor used by dendritic

cells and thymic epithelial cells to direct captured antigens from the extracellular space to specialized antigen processing. It mediates antigen uptake and presentation and cross-presentation to T cells. It has been reported that CD205 acts as a recognition receptor for dying cells, potentially provides an important pathway for the uptake of self-antigen in the intrathymic environment, and is involved in peripheral tolerance. Antibody-mediated antigen-targeting via the DEC-205 receptor increases the efficiency of vaccination for T cell immunity.

**Antigen
References:**

1. Jiang WP, *et al.* 1995. *Nature* 375:151.
2. Small M and Kraal G. 2003. *Int. Immunol.* 15:197.
3. Shrimpton RE, *et al.* 2009. *Mol. Immunol.* 46:1229.