

PE anti-mouse TCR Vβ5.1, 5.2

Catalog # / Size: 1297520 / 100 µg
1297515 / 25 µg

Clone: MR9-4

Isotype: Mouse IgG1, κ


Immunogen: Murine T cell hybridoma 2HB51.8

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2




C57BL/6 splenocytes stained with CD3e (145-2C11) APC and MR9-4 PE

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



C57BL/6 splenocytes stained with CD3e (145-2C11) APC and mouse IgG1 PE isotype control

Application Notes: Additional reported applications (for the relevant formats) include: Induction of proliferation of Vβ5.1⁺ and Vβ5.2⁺ T cells^{2, 3} and *in vivo* depletion of Vβ5⁺ T cells⁴.

- Application References:**
1. Kanagawa O, *et al.* 1991. *J. Immunol.* 147:1307. (FC)
 2. Kanagawa O, *et al.* 1992. *J. Immunol.* 149:9. (Activ)
 3. Woodland DL, *et al.* 1993. *J. Exp. Med.* 177:433. (Activ)
 4. Gelber C, *et al.* 1992. *Cancer Res.* 52:6507. (Deplete)

Description: Vβ5.1 and 5.2 T cell receptor (TCR Vβ5.1, 5.2) are variants of TCR β chain that, along with TCR α chain, forms the TCR heterodimer. In association with the CD3 complex, TCR α/β is responsible for antigen recognition in the MHC-Peptide complex and the initiation of T cell-mediated immune responses.

- Antigen References:**
1. Marrack P, *et al.* 2008. *Annu. Rev. Immunol.* 26:171.
 2. Sim GK and Augustin AA. 1985. *Cell* 42:89.
 3. Mami-Chouaib F, *et al.* 2002. *Immunol. Rev.* 188:114.