
Product Data Sheet

PE/Cy7 anti-mouse CD138 (Syndecan-1)

Catalog # / Size:	1312565 / 25 μg 1312570 / 100 μg
Clone:	281-2
Isotype:	Rat IgG2a, к
Immunogen:	Mouse mammary gland epithelial cell line NMuMG
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	0.2



C57BL/6 mouse splenocytes were stained with CD19 APC and CD138 (clone 281-2) PE/Cy7 (top) or rat IgG2a, κ PE/Cy7 isotype control (bottom).

105

Applications:

		10" -
Applications:	Flow Cytometry	8
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.125 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.	Rat IgG2a, k PE/Cy
Application Notes:	Additional reported applications (for the relevant formats) include: immunohistochemical staining of frozen tissue3 and formalin-fixed paraffin embedded tissue ⁴ and immunofluorescent staining ^{2,3} .	CD19 APC
Application References:	 Jalkanen M, <i>et al.</i> 1985. <i>J. Cell. Biol.</i> 101: Miettinen H, <i>et al.</i> 1994. <i>J. Cell. Sci.</i> 107: Li Q, <i>et al.</i> 2002. <i>Cell</i> 111:635. (IF, IHC) McCarthy BA, <i>et al.</i> 2012. <i>BMC Cancer.</i> 1 	1571. (IF)
Description:	CD138, a member of the syndecan protein heparin sulfate proteoglycan also known a	

Description: CD138, a member of the syndecan protein family, is a type I integral membrane heparin sulfate proteoglycan also known as Syndecan-1. Syndecan-1 participates in cell proliferation, cell migration, and cell matrix adhesion via interaction with collagen, fibronectin, and other soluble molecules (such as FGF-basic). It is expressed on normal and malignant plasma cells, pre-B cells, mesenchymal cells, epithelial cells, and endothelial cells.

Antigen	1. Zong F, <i>et al.</i> 2011. <i>PLoS ONE</i> 6:e14816.	
---------	--	--

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com
 References:
 2. Yamashita Y, et al. 1999. J. Immunol. 162:5940.

 3. Sanderson RD, et al. 1989. Cell. Regul. 1:27.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com