


**PE/Dazzle™ 594 anti-mouse CD138 (Syndecan-1)**

<b>Catalog # / Size:</b>	1312640 / 100 µg 1312635 / 25 µg
<b>Clone:</b>	281-2
<b>Isotype:</b>	Rat IgG2a, κ
<b>Immunogen:</b>	Mouse mammary gland epithelial cell line NMuMG
<b>Reactivity:</b>	Mouse
<b>Preparation:</b>	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.2



C57BL/6 mouse bone marrow cells were stained with CD19 APC and CD138 (clone 281-2) PE/Dazzle™ 594 (top) or rat IgG2a, κ PE/Dazzle™ 594 isotype control (bottom). Data shown is gated on lymphoid cell population.


**Applications:**

<b>Applications:</b>	Flow Cytometry
<b>Recommended Usage:</b>	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

<b>Application Notes:</b>	Additional reported applications (for the relevant formats) include: immunohistochemical staining of frozen tissue <sup>3</sup> and formalin-fixed paraffin embedded tissue <sup>4</sup> and immunofluorescent staining <sup>2,3</sup> .
---------------------------	--

<b>Application References:</b>	1. Jalkanen M, <i>et al.</i> 1985. <i>J. Cell. Biol.</i> 101:976. (FC) 2. Miettinen H, <i>et al.</i> 1994. <i>J. Cell. Sci.</i> 107:1571. (IF) 3. Li Q, <i>et al.</i> 2002. <i>Cell</i> 111:635. (IF, IHC) 4. McCarthy BA, <i>et al.</i> 2012. <i>BMC Cancer.</i> 12:203. (IHC)
--------------------------------	--



**Description:** CD138, a member of the syndecan protein family, is a type I integral membrane

heparin sulfate proteoglycan also known as Syndecan-1. Syndecan-1 participates in cell proliferation, cell migration, and cell matrix adhesion via interaction with collagen, fibronectin, and other soluble molecules (such as FGF-basic). It is expressed on normal and malignant plasma cells, pre-B cells, mesenchymal cells, epithelial cells, and endothelial cells.

**Antigen  
References:**

1. Zong F, *et al.* 2011. *PLoS ONE* 6:e14816.
2. Yamashita Y, *et al.* 1999. *J. Immunol.* 162:5940.
3. Sanderson RD, *et al.* 1989. *Cell. Regul.* 1:27.