

PE/Cy7 anti-mouse PD-1H (VISTA)

Catalog # / Size: 1318570 / 100 µg
1318565 / 25 µg

Clone: MH5A

Isotype: Hamster IgG


Immunogen: PD-1H- IgG Fc fusion protein

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2




C57BL/6 mouse splenocytes were stained with CD8 FITC and PD-1H (clone MH5A) PE/Cy7 (top) or Armenian Hamster IgG PE/Cy7 isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of graft vs host disease (GVHD), Western blotting, and immunohistochemical staining of paraffin embedded tissue sections.



Application References: 1. Flies DB, *et al.* 2011. *J. Immunol.* 187:1537.

Description: PD-1H, also known as VISTA, is a 309 aa type I transmembrane protein, composed of seven exons. PD-1H has one Ig-V like domain, and its sequence is similar to the Ig-V domains of the members of CD28 and B7 families. PD-1H is expressed by a subset of T cells, macrophages, dendritic cells, neutrophils, and NK cells. It has been proposed that PD-1H can be useful to modulate the host immune response to allogeneic transplants.

Antigen References: 1. Flies DB, *et al.* 2011. *J. Immunol.* 187:1537.
2. Wang Li, *et al.* 2011. *J. Exp Med.* 208:577.