


Purified anti-mouse CD185 (CXCR5)

Catalog # / Size: 1327510 / 100 µg
Clone: L138D7
Isotype: Rat IgG2b, κ
Immunogen: mCXCR5-transfected cells
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



C57BL/6 mouse splenocytes were stained with purified CXCR5 (clone L138D7, filled histogram) or rat IgG2b, κ isotype control (open histogram), followed by anti-rat IgG PE.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone L138D7 staining works optimally at room temperature or 4°C. Unlike other chemokine receptor antibodies, avoid using L138D7 at 37°C.

Description: CD185 is also known as CXCR5. It is the receptor for chemokine CXCL13/BLC, which is chemotactic for B cells. CXCR5 is expressed on B cells and a subset of T cells in the spleen, neuronal tissue, lymph nodes, and bone marrow. It is important for migration of B cells into the B cell follicles of the spleen and Peyer's patches. Follicular helper T cells (Tfh) also express CXCR5 and the ability of these cells to migrate to the lymph node is modulated by the balanced expression of CCR7 and CXCR5.

Antigen References:

1. Kaiser E, *et al.* 1993. *Eur. J. Immunol.* 23:2532.
2. Forster R, *et al.* 1994. *Cell. Mol. Biol.* 40:381.
3. Forster R, *et al.* 1994. *Blood* 84:830.
4. Forster R, *et al.* 1996.