


Brilliant Violet 785™ anti-mouse CD185 (CXCR5)


Catalog # / Size:	1327615 / 50 µg
Clone:	L138D7
Isotype:	Rat IgG2b, κ
Immunogen:	mCXCR5-transfected cells
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 785™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 785™ and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Concentration:	NULL



C57BL/6 mouse splenocytes were stained with CD45R/B220 PE and CXCR5 (clone L138D7) Brilliant Violet 785™ (top) or rat IgG2b, κ Brilliant Violet 785™ isotype control (bottom).

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	Brilliant Violet 785™ excites at 405 nm and emits at 785 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 785™ is a trademark of Sirigen Group Ltd.
Application Notes:	Clone L138D7 staining works optimally at room temperature or 4°C. Unlike other chemokine receptor antibodies, avoid using L138D7 at 37°C.



Description: CD185 is also known as CXCR5. It is the receptor for chemokine CXCL13/BLC,

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

which is chemotactic for B cells. CXCR5 is expressed on B cells and a subset of T cells in the spleen, neuronal tissue, lymph nodes, and bone marrow. It is important for migration of B cells into the B cell follicles of the spleen and Peyer's patches. Follicular helper T cells (Tfh) also express CXCR5 and the ability of these cells to migrate to the lymph node is modulated by the balanced expression of CCR7 and CXCR5.

**Antigen
References:**

1. Kaiser E, *et al.* 1993. *Eur. J. Immunol.* 23:2532.
2. Forster R, *et al.* 1994. *Cell. Mol. Biol.* 40:381.
3. Forster R, *et al.* 1994. *Blood* 84:830.
4. Forster R, *et al.* 1996.