

PE/Dazzle™ 594 anti-mouse CD184 (CXCR4)

Catalog # / Size: 1332570 / 100 µg
1332565 / 25 µg

Clone: L276F12

Isotype: Rat IgG2b, κ


Immunogen: Mouse CXCR4-transfected cells

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse thymocytes were stained with CD4 FITC and CD184 (clone L276F12) PE/Dazzle™ 594.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Description: CD184, also known as CXCR4, is a member of the G protein coupled receptor family that binds CXCL12 (SDF1). CXCR4 and CXCL12 play an important role in immune and inflammatory responses through the regulation of cell migration and growth. CXCR4 plays a crucial role in the pathogenesis of several autoimmune diseases such as atherosclerosis, rheumatoid arthritis, and wound healing. In addition, CXCR4 is the cofactor for fusion and entry of the T cell-tropic form of HIV-1.

- Antigen References:**
1. Kucia M, *et al.* 2005. *Stem Cells* 23:879.
 2. Muller A, *et al.* 2001. *Nature* 410:50.
 3. Saini V, *et al.* 2010. *J. Biol. Chem.* 285:15566.
 4. Prasad A, *et al.* 2007. *J. Leuko*