

APC anti-mouse P2X7R

Catalog # / Size: 1343525 / 25 µg
1343530 / 100 µg

Clone: 1F11

Isotype: Rat IgG2b, κ


Immunogen: Murine colon mast cells

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: Lot-specific




C57BL/6 splenocytes were stained with CD4 Pacific Blue™ and P2X7R (clone 1F11) APC (top) or rat IgG2b, κ APC isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications for the relevant formats include: immunoprecipitation¹, Western blotting¹, immunohistochemistry¹, *in vivo* inhibition of intestinal inflammation and mast cell activation¹.

Application References: 1. Kurashima Y, *et al.* 2012. *Nat. Commun.* 3:1034. (FC, IP, WB, IHC, FA)

Description: P2X7R, also known as P2X7 receptor, belongs to the family of ligand-gated ion channel receptors. It is expressed on T cells, B cells, macrophages, and microglia. The receptor opens in the presence of extracellular ATP or NAD, leading to intracellular calcium mobilization. P2X7R activation requires higher concentrations of ATP compared to other P2X receptors. Longer stimulation results in larger pores, allowing passage of larger molecules. Activation of these molecules also leads to mitochondrial and cytoskeletal changes as well as IL-1 β maturation and release. Ligation of P2X7 receptor can lead to membrane blebbing and cell death.

Antigen
References:

1. Surprenant A, *et al.* 1996. *Science*. 272:735.
2. Chused TM, *et al.* 1996. *J. Immunol.* 157:1371.
3. Gargett CE, *et al.* 1997. *Br. J. Pharmacol.* 122:911.
4. Kawamura H, *et a*