
Product Data Sheet

Alexa Fluor[®] 488 anti-mouse CX3CR1


| Catalog # / Size: | 1345110 / 100 μg 1345105 / 25 μg |
|--------------------|---|
| Clone: | SA011F11 |
| Isotype: | Mouse IgG2a, к |
| Immunogen: | Mouse CX3CR1-transfected cells |
| Reactivity: | Mouse |
| Preparation: | The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 488 under optimal conditions. |
| Formulation: | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. |
| Concentration: | 0.5 |



C57BL/6 mouse splenocytes were stained with CD11b APC and CX3CR1 (clone SA011F11) Alexa Fluor® 488 (top) or mouse IgG2a, κ Alexa Fluor® 488 isotype control (bottom).

Applications:

| Applications: | Flow Cytometry |
|-----------------------|---|
| Recommended Usage: | Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.06 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application. |
| | * Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm. |
| Application Notes: | For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF [™] purified antibody (Cat. No. 149011) with a lower endotoxin limit than standard LEAF [™] purified antibodies (Endotoxin <0.01 EU/microg). |
| | |



Description: CX3CR1 is a 40 kD, G-protein coupled receptor, with seven transmembrane regions. CX3CR1 is expressed by resident and alternatively activated macrophages (M2), a subset of monocytes, dendritic cells (DCs), NK cells, a subset of memory T cells, and mast cells. CX3CR1 is involved in cell recruitment during inflammation and participates in cell adhesion and extravasation from blood vessels. Its ligand is CX3CL1, also known as fractalkine or neurotactin. CX3CR1 is also a coreceptor for HIV1 and variations in this gene leads to increased susceptibility to HIV. In the brain, it is expressed by glial cells, which interact with CX3CL1 expressed by neurons.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

| Antigen | 1. Ponzetta A, <i>et al.</i> 2013. <i>J. Immunol.</i> 191:5684. |
|--------------------|---|
| References: | 2. Jacquelin S, <i>et al.</i> 2013. <i>Blood.</i> 122:674. |
| | 3. Garcia JA, <i>et al.</i> 2013. <i>J. Immunol.</i> 191:1063. |
| | 4. Lee YS, <i>et al.</i> 2013. <i>Ce</i> |

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com