Product Data Sheet

APC/Cy7 anti-mouse CD20

Catalog # / Size: 1352090 / 100 µg

1352085 / 25 µg

Clone: SA275A11 Isotype: Rat IgG2b, ĸ

Mouse CD20 - transfected cells Immunogen:

Reactivity: Mouse

Preparation: The antibody was purified by affinity

chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7


and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

each application.

containing 0.09% sodium azide.

Concentration: 0.2 mg/ml




Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 μα per million cells in 100 μl volume. It is recommended that the reagent be titrated for optimal performance for



C57BL/6 mouse splenocytes were stained with CD45R/B220 FITC and CD20 (clone SA275A11) APC/Cy7 (left) or rat IgG2b, κ APC/Cy7 isotype control (right).

Application References:

- 1. Morsy DE, et al. 2013. J. Immunol. 191:3112.
- 2. Lund FE. Randall TD. 2010. Nat. Rev. Immunol. 10:236.
- 3. Beers SA, et al. 2010. Blood 115:5191.
- 4. Kuijpers TW, et al. 2010. /.

Description:

CD20 is a 33-37 kD protein, a member of the MS4A family, with four transmembrane spanning regions that present as a homo-oligomeric complexes in the cell surface when associating with MHC class I and II, CD53, CD81, and CD82. CD20 is expressed on B cells and a subset of T cells, but not on plasma cells. CD20 regulates B-cell activation and proliferation. Its ligation promotes transmembrane Ca²⁺ trafficking. CD20 is an important therapeutic target in the treatment of B cell lymphomas and leukemias.

Antigen References:

- 1. Morsy DE, et al. 2013. J. Immunol. 191:3112.
- 2. Lund FE. Randall TD. 2010. Nat. Rev. Immunol. 10:236.
- 3. Beers SA, et al. 2010. Blood 115:5191.

4. Kuijpers TW, <i>et al.</i> 2010. <i>J.</i>