


Purified anti-mouse CD186 (CXCR6)


Catalog # / Size: 1355510 / 100 µg
Clone: SA051D1
Isotype: Rat IgG2b, κ
Immunogen: mCXCR6-transfectants
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.2



C57BL/6 mouse splenocytes were stained with CD3 APC and purified CD186 (clone SA051D1, top) or purified rat IgG2b, κ isotype control (bottom), followed by anti-rat IgG PE.

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Description: CD186, or CXCR6, is a 39 kD G-protein coupled chemokine receptor with seven transmembrane-spanning regions. Its ligand is CXCL16. It is expressed on different T cell subsets and is upregulated in activated T cells. Expression of CXCR6 is correlated with increased inflammatory responses and seems to contribute to liver fibrosis.

- Antigen**
- References:**
1. Kim CH, *et al.* 2001. *J. Clin. Invest.* 107:595.
 2. Heesch K, *et al.* 2014. *PLoS One.* 9:5.
 3. Wehr A, *et al.* 2013. *J. Immunol.* 190:5226.