


**KIRAVIA Blue 520™ anti-mouse MERTK (Mer)****Catalog # /** 1357640 / 100 µg**Size:** 1357635 / 25 µg**Clone:** 2B10C42**Isotype:** Rat IgG2a, κ**Immunogen:** Mouse MERTK extracellular domain**Reactivity:** Mouse**Preparation:** The antibody was purified by affinity chromatography and conjugated with KIRAVIA Blue 520™ under optimal conditions.**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide**Concentration:** 0.2 mg/mL

C57BL/6 mouse splenocytes were stained with CX3CR1 APC and anti-mouse MERTK (Mer) (clone 2B10C42) KIRAVIA Blue 520™ (left) or rat IgG2a, κ KIRAVIA Blue 520™ isotype control (right).

**Applications:****Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* KIRAVIA Blue 520™ has an excitation maximum of 495 nm, and a maximum emission of 520 nm.

**Description:** MerTK (Mer) is a member of the TAM (TYRO3/AXL/MerTK) family. It is a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type domains, and one tyrosine kinase domain. MerTK is mainly expressed by macrophages, monocytes, and dendritic cells. Its ligands are LGALS3, TUB, TULP1, and GAS6. MerTK is involved in the regulation of TLR signaling, efferocytosis, phagocytosis, cell survival, macrophage migration, and the inhibition of inflammation.

**Antigen**  
**References:**

1. Zagórska A, *et al.* 2014. *Nat. Immunol.* 15:920.
2. Toda S, *et al.* 2014. *Blood* 123:3963.
3. Chung WS, *et al.* 2013. *Nature* 504:394.
4. Carrera Silva EA, *et al.* 2013. *Immunity* 39:160.
5. Yi Z, *et al.* 2009. *Blood* 114:3191.