

APC anti-mouse CD95 (Fas)

Catalog # / Size: 1363015 / 25 µg
1363020 / 100 µg

Clone: SA367H8

Isotype: Mouse IgG1, κ


Immunogen: Mouse Fas Transfectants

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: Lot-specific



C57BL/6 thymocytes were stained with CD95 (clone SA367H8) APC (filled histogram) or mouse IgG1, κ APC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.125 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Description: CD95, also known as Fas, is an approximately 45 kD type I transmembrane protein belonging to the TNFR superfamily (TNFRSF6). The expression of CD95 has been described in various organs, such as thymus, spleen, liver, heart, lung and ovary. Upon ligand (FasL) binding, CD95 forms the death-inducing signaling complex (DISC) intracellularly and induce apoptosis of the cell. CD95-induced apoptosis plays important roles in development, as well as in maintaining peripheral tolerance of the immune system.

Antigen
References:

1. Ogasawara J, *et al.* 1995. *J. Exp. Med.* 181:485.
2. Ogasawara J, *et al.* 1993. *Nature.* 364:806.
3. Nishimura Y, *et al.* 1995. *J. Immunol.* 154:4395.