PerCP/Cyanine5.5 anti-mouse CD204

Catalog # / Size:	1373575 / 25 μg 1373580 / 100 μg	
Clone:	1F8C33	
lsotype:	Rat IgG2a	
Immunogen:	Recombinant mouse CD204 extracellular domain	
Reactivity:	Mouse	
Preparation:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide	ſ
Concentration:	0.2 mg/mL	r V



Mouse leukemic monocytemacrophage cell line RAW267.4 was stained with PerCP/Cyanine5.5 anti-mouse CD204 (clone 1F8C33) (filled histogram) or rat IgG2a, ĸ PerCP/Cyanine5.5 isotype control (open histogram).

Applications:

Applications:	Flow Cytometry
---------------	----------------

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5 \ \mu$ g per million cells in 100 μ L volume. It is recommended that the reagent be titrated for optimal performance for each application.

 \ast PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

ApplicationThis clone has minimal recognition of CD204 in C57BL/6.Notes:

Description: CD204, also known as scavenger receptor A (SR-A) and the macrophage scavenger receptor (MSR), is 220 kDa, trimeric type II transmembrane protein, with one scavenger receptor cysteine-rich domain (SRCR). It is a phagocytic pattern-recognition receptor (PRR) expressed on macrophages and dendritic cells. CD204 is a receptor mediating recognition and internalization of low-density lipoprotein (LDL) by macrophages and plays a critical role in atherogenesis. CD204 also recognizes apoptotic cells, modified lipid proteins, and exogenous pathogen-associated molecular patterns (PAMPs), which results in the induction of innate immune and inflammatory responses. CD204 can act as a co-receptor for Toll-like receptors, such as TLR3, TLR4, or TLR9, to facilitate the expression of proinflammatory cytokines. CD204 has been implicated in several pathological processes such as Alzheimer's disease, sepsis, ischemic injury, and coronary artery disease.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Antigen **References:**

- 1. Shichita T, et al. 2017. Nat Med. 23:723.
- Muczynski V, et al. 2017. Blood. 129:2443.
 Iftakhar-E-Khuda I, et al. 2016. Proc Natl Acad Sci U S A. 113:10643.
 Nellimarla S, et al. 2015. J Immunol. 195:3858.
- 5. Bonilla DL, et al. 2013. Immunity. 39:537.