

PE/Dazzle™ 594 anti-mouse CD183 (CXCR3)

Catalog # / Size: 1379565 / 25 µg
1379570 / 100 µg

Clone: S18001A

Isotype: Rat IgG2b, κ


Immunogen: Mouse CXCR3-transfectants

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Concentration: 0.2 mg/mL



C57BL/6 splenocytes were stained with CD3ε FITC and CD183 (CXCR3) (clone S18001A) PE/Dazzle™ 594 (left) or rat IgG2b, κ PE/Dazzle™ 594 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Description: CD183, also known as CXCR3, is a member of the C-X-C chemokine family, characterized by a pair of cysteine residues separated by a single amino acid. CXCR3 is a 38 kD seven pass transmembrane receptor coupled to G-protein. It mediates Ca²⁺ mobilization and chemotaxis in response to C-X-C chemokines, such as IP10 (CXCL10), MIG (CXCL9), I-TAC (CXCL11) and PF4 (CXCL4). CXCR3 is expressed primarily on activated T lymphocytes, NK cells, and some epithelial cells and endothelial cells. It is not expressed on B cells, monocytes, or granulocytes.

- Antigen References:**
1. Farber JM. 1997. *J. Leukoc. Biol.* 61(3):\246-57.
 2. Bonecchi R, et al. 1998. *J. Exp. Med.* 187(1):129-34.
 3. Aota K, et al. 2018. *J. Oral. Pathol. Med.* 12756.
 4. Kim B, et al. 2018. *Data. Brief.* 18:518-522.
 5. Saahene RO, et al. 2018. *Cancer Biother. Radiopharm.* 2450.