


Biotin anti-rat CD11b/c

Catalog # / Size: 1609015 / 100 µg
Clone: OX-42
Isotype: Mouse IgG2a, κ
Immunogen: Rat peritoneal macrophages.
Reactivity: Rat
Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5




LOU rat bone marrow cells stained with biotinylated OX-42, followed by Sav-PE (gated on myeloid cells)

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemistry of acetone-fixed frozen sections^{1,2}, immunoprecipitation³, *in vivo* and *in vitro* blocking of C3bi binding^{3,4}.

- Application References:**
- Whiteland JL, *et al.* 1995. *J. Histochem. Cytochem.* 43:313. (IHC)
 - Milligan CE, *et al.* 1991. *J. Comp. Neurol.* 314:125. (IHC)
 - Robinson AP, *et al.* 1986. *Immunology* 57:239. (Block)
 - Issekutz SE, *et al.* 1992. *Immunology* 76:655. (Block)
 - Muehlbauer SM, *et al.* Am. J Pathol. 177:735. (FC) [PubMed](#)
 - Thomas RA, *et al.* 2009. *J Pharmacol Toxicol Methods.* 60:263. [PubMed](#)
 - Takeda Y, *et al.* 2011. *Immunobiology.* 216:1094. [PubMed](#)
 - Furuhashi K, *et al.* 2013. *J Am Soc Nephrol.* [PubMed](#)
 - Cheng HY, *et al.* 2013. *Cytotherapy.* 1465:608. [PubMed](#)



LOU rat splenocytes stained with biotinylated OX-42, followed by Sav-PE

Description: The OX-42 antibody reacts with the CR3 complement (C3bi) receptor expressed on monocytes, granulocytes, macrophages, dendritic cells, NK cells, and a subset of lymphocytes. This antibody appears to recognize a common epitope shared between CD11b and CD11c (integrin α_M and α_X chains). The OX-42 antibody precipitates three polypeptides with apparent molecular weights of 160, 103, and

95 kD, respectively. This antibody has been shown to block the formation of complement-mediated rosettes and leukocyte migration.

- Antigen** 1. Robinson AP, *et al.* 1986. *Immunology* 57:239.
References: 2. Barcaly AN. 1981 *Immunology* 42:593.