


APC anti-rat CD47**Catalog # /** 1620570 / 100 µg**Size:** 1620565 / 25 µg**Clone:** OX-101**Isotype:** Mouse IgG1, κ**Immunogen:** Mouse CXCR3-transfectants**Reactivity:** Rat**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.**Concentration:** 0.2 mg/ml

LOU rat splenocytes were surface stained with CD47 (clone OX-101) APC (filled histogram)

Applications:**Applications:** Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application

- References:**
1. Vernon-Wilson, *et al.* 2000. *Eur. J. Immunol.* 30:2130.
 2. Brown E, *et al.* 1990. *J. Cell. Biol.* 111:2785.
 3. Mawby WJ, *et al.* 1994. *Biochem. J.* 304:525.

Description: CD47, also known as Integrin-Associated Protein (IAP), is a membrane protein of about 50 kD with an IgV-like extracellular domain, a five membrane-spanning segment, and a short terminal cytoplasmic region. It is widely expressed on many cell types and often associated with beta 3 integrins. The significance of this molecule is recently drawing increasing attention. It has been reported that CD47 functions as a self marker. Red cells lacking CD47 were rapidly cleared from the bloodstream by splenic macrophages. By binding to SIRPα, CD47 controls hemostatic innate immune functions, such as phagocytosis and cell trafficking.

Antigen

- References:**
1. Vernon-Wilson, *et al.* 2000. *Eur. J. Immunol.* 30:2130.
 2. Brown E, *et al.* 1990. *J. Cell. Biol.* 111:2785.
 3. Mawby WJ, *et al.* 1994. *Biochem. J.* 304:525.